

Humidifier Bacteria-Algae Treatment

Humidifier Bacteria-Algae treatment is a bacteriostat which can control the build-up of bacteria and algae in water tanks of portable, (evaporative) humidifiers.

ACTIVE INGREDIENTS

n-Alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ 5% C ₁₈) dimethyl benzyl ammonium chlorides.....	2.25%
n-Alkyl (68% C ₁₂ , 32% C ₁₄) dimethyl ethylbenzyl ammonium chlorides.....	2.25%
INERT INGREDIENTS.....	95.50%
Total	100.00%

KEEP OUT OF REACH OF CHILDREN

WARNING

See side panels for additional precautionary statements.

EPA REG. NO. 1839-120
EPA EST. NO. 1839-IL-01

STEPAN COMPANY
22 West Frontage Road • Northfield, IL 60093

NET CONTENTS

(center panel)

Approved 06/03/10

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Humidifier Bacteria-Algae Treatment is designed for use in portable, (evaporative) humidifiers.

Be sure humidifier water tank and filters are thoroughly cleaned before each heating season.

For every gallon of water contained in the humidifier tank, add (one capful) (1/6 of an ounce) (or 1 ounce per 6 gallons) of this product. When you refill the tank, add (one capful) (1/6 of an ounce) (or 1 ounce per 6 gallons) of this product for every gallon of water added.

Not for heat vaporizing or atomizing type vaporizer humidifiers.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS.

WARNING

KEEP OUT OF REACH OF CHILDREN. Causes substantial but temporary eye injury and skin irritation. Do not get in eyes, on skin or on clothing. Wear safety glasses. Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

(OPTIONAL) You may contact xxx/xxx-xxxx for emergency medical treatment information.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage.

STORAGE AND DISPOSAL

PESTICIDE STORAGE - Store in a dry place no lower in temperature than 50°F or higher than 120°F. Store in original container in a locked storage area inaccessible to children.

PESTICIDE DISPOSAL - Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office.

(For Industrial, Institutional, Commercial Products)

CONTAINER DISPOSAL – Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Triple rinse container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after flow begins to drip. Fill container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

(For Household/Residential Products)

CONTAINER DISPOSAL - Nonrefillable container. Do not reuse or refill this container. Wrap container and discard in trash. Offer for recycling, if available.